

ISWAVLD 2⁽¹⁾23

International Symposium of the World Association of Veterinary Laboratory Diagnosticians 29 JUNE-1 JULY 2023 Congress Centre

Lyon

Towards the veterinary diagnostics of the future

Main topic : Antimicrobial resistance: A worldwide concern

Evaluation of the antimicrobial effect of Tocosh (Solanum tuberosum L.) on Campylobacter spp. isolated from chickens in markets of Lima, Peru: Preliminary results

CORTEZ RODRIGUEZ C. ¹, DÍAZ COAHILA D. ¹, ANGULO HERRERA P. ¹, CONTE JUNIOR C. ², AGUILAR FELICES E. ³, ENCISO ROCA E. ³, TINCO JAYO J. ³, ROJAS MONTES M. ⁴, <u>LAZARO DE LA TORRE C. ¹</u>

¹ Laboratorio de Farmacología y Toxicología Veterinaria, Facultad de Medicina Veterinaria, Universidad Nacional Mayor de San Marcos, Lima, Peru; ² Instituto de Química, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil; ³ Departamento Académico de Medicina Humana, Facultad de Ciencias de la Salud, Universidad Nacional de San Cristóbal de Huamanga, Ayacucho, Peru; ⁴ Laboratorio de Inmunología y Virología Veterinaria, Facultad de Medicina Veterinaria, Universidad Nacional Mayor de San Marcos, Lima, Peru

Introduction: Tocosh is a traditional food based on potato (*Solanum tuberosum* L.) obtained by fermentation, and the Peruvian Andean populations use this product for its antibiotic capacity. Its effect against pathogenic bacteria such as *E. coli, S. aureus, L. monocytogenes*, and *S. mutans* has been demonstrated experimentally and even in strains resistant to conventional antibiotics. *Campylobacter* spp. is the fourth bacterial pathogen that causes human gastrointestinal disease. This microorganism, found in the broiler's gut, has increased its antibiotic resistance. This work aims to evaluate the antimicrobial effect of Tocosh flour on *Campylobacter* spp. isolated to chicken meat from markets in Lima, Peru. **Methods:** For this purpose, 35% alcoholic solutions for three commercial Tocosh flours (T1, T2, and T3) were prepared and left to macerate for seven days. The product was filtered with Whatman N°1 paper and concentrated with a rotary evaporator (60°C) up to 5% of the original volume. Concentrations of 100, 75, 50, 50, 25, and 0% were prepared from this solution. The agar well-diffusion method was performed with 0.5 on the McFarland scale (600OD) of *Campylobacter coli* isolated from chicken meat, *C. coli* (ATCC 33559), *Escherichia coli* (ATCC 25922), and *Staphylococcus aureus* (ATCC 29213). Bacteria solutions were seeded on plates with Muller-Hinton blood agar (5%); then wells were made in the agar to deposit 35 µL of the different Tocosh solutions (100, 75, 50, 50, 25, 0%). Plates were incubated at 42°C/48h in microaerophilic. **Results:** The results show that the T1 flour had an inhibition zone of 9 and 8 mm at 100% and 75% for the *C. coli* strain isolated from chickens in the 4TCC strain, only the 100% concentration presented an inhibition zone of 8 mm. The other two flours (T2 and T3) showed color changes in the agar as an inhibition zone but with colony growth and were therefore discarded. The 100% concentrations of the three flour brands achieved inhibition zones of 9, 8, and 7 mm were obse

Funding: This study was supported by Universidad Nacional Mayor de San Marcos within the framework of the call PCONFIGI (Code: A22081111) and PINTERDIS (Code A2108001i)